

# Trends in extreme precipitation indices derived from a daily rainfall database for the South of Portugal

Ana Cristina Costa<sup>a\*</sup> and Amílcar Soares<sup>b</sup>

<sup>a</sup> ISEGI, Instituto Superior de Estatística e Gestão de Informação, Universidade Nova de Lisboa, Portugal

<sup>b</sup> CERENA, Centro de Recursos Naturais e Ambiente, Instituto Superior Técnico, Portugal

**ABSTRACT:** The rainfall regime of the South of Portugal is Mediterranean with Atlantic influence. Long-term series of *reliable* precipitation records are essential for land and water resources management, climate-change monitoring, modelling of erosion and run-off, among other applications for ecosystem and hydrological impact modelling. This study provides a qualitative classification of 106 daily rainfall series from stations located in the South of Portugal and evaluates temporal patterns in extreme precipitation by calculating a number of indicators at stations with homogeneous data within the 1955/1999 period. The methodology includes both absolute and relative approaches and a new homogeneity testing procedure, besides the application of other statistical tests. The proposed technique is an extension of the Ellipse test that takes into account the contemporaneous relationship between several candidate series from the same climatic area (*SUR+Ellipse test*). The results indicate that this technique is a valuable tool for the detection of non-climatic irregularities in climate time series if the station network is dense enough. The existence of trends and other temporal patterns in extreme precipitation indices was investigated and uncertainty about rainfall patterns evolution was assessed. Three indices describing wet events and another three indicators characterizing dry conditions were analysed through regression models and smoothing techniques. The simple aridity intensity index (AII) reflects increases in the magnitude of dryness. Especially pronounced trends are found over most of southern Portugal in the 1955/1999 period, highlighting the fact that large areas are threatened by drought and desertification. The trend signals of the wetness indices are not significant at the majority of stations, but there is evidence of increasing short-term precipitation intensity over the region during the last three decades of the twentieth century. Finally, the results also indicate that extreme precipitation variability and climate uncertainty are greater in recent times. Copyright © 2008 Royal Meteorological Society

KEY WORDS aridity; climate variability; extreme precipitation; homogeneity testing; Portugal; rainfall intensity; trend analysis

Received 5 January 2008; Revised 6 November 2008; Accepted 11 November 2008

## 1. Introduction

Portugal is geographically located in the southwesterly extreme of the Iberian Peninsula (between 37° and 42°N and 6.5° and 9.5°W). Global circulation and regional climatic factors (e.g. latitude, orography, oceanic and continental influences) explain the spatial distribution of rainfall, as well as its intra-annual variability, i.e. seasonal variability (Trigo and DaCamara, 2000; Goodess and Jones, 2002). The precipitation regimes are of a different nature in northern and southern regions of Portugal: in the North the precipitation regime has an orographic origin, whereas in the South it is associated to cyclogenetic activity (Trigo and DaCamara, 2000). The inter-annual variability is of a different nature, since the circulation variability is insufficient to explain the observed inter-annual variability of rainfall (Trigo and DaCamara, 2000; Goodess and Jones, 2002; Haylock and Goodess, 2004). In southern Portugal, summer precipitation, almost close to zero during this season, is sometimes associated with

local convective activity. These storms can occur with a large degree of independence from the circulation weather type, which characterizes the Iberian circulation for that specific day (Trigo and DaCamara, 2000).

Recent studies, based on climate models and past observed records, predict a future increase in droughts in the South of Europe as a result of increased evapotranspiration and a relatively slow decrease of rainfall amounts and precipitation frequency (e.g. Kostopoulou and Jones, 2005; Vicente-Serrano and Cuadrat-Prats, 2007). The results obtained by Goodess and Jones (2002) for the Portuguese stations show general agreement with those from Trigo and DaCamara (2000) who considered ten classes of weather circulation types for Portugal. Their results suggest that the cyclonic class is associated with a fairly homogeneous distribution of precipitation over most of the country. Moreover, the 'rainy' classes with an Atlantic origin (mainly W and SW; NW to a lesser degree) are to be associated with the observed strong decrease in precipitation from North to South.

In arid and semi-arid regions such as the South of continental Portugal, research on the extent of dryness and temporal trends in heavy rainfall events is an important

\* Correspondence to: Ana Cristina Costa, ISEGI, Universidade Nova de Lisboa, Campus de Campolide, 1070-312 Lisbon, Portugal.  
E-mail: ccosta@isegi.unl.pt

contribution to evaluate desertification dynamics and to identify areas potentially at risk from land degradation. However, studies focussing on the role of regional climate change on erosivity and aridity factors are lacking for this region, especially at the local scale. This study attempts to compile a daily rainfall database for the South of Portugal and subsequently to evaluate temporal trends in extreme precipitation by calculating a number of climate indices.

Besides the description of the study domain and precipitation data, the first part of the article focuses on the detection of temporal discontinuities in the precipitation time series. This issue is of major importance, because non-climatic factors make data unrepresentative of the actual climate variation and might bias the studies' conclusions. A break could result from a recalibration of an instrument or a station relocation; a linear trend could result from a gradual but constant degradation of a sensor; and a non-linear trend could result from vegetative growth around the instruments. Several techniques have been developed for detecting inhomogeneities in time series of weather elements. The approaches underlying the homogenization techniques are quite different and typically depend on the type of element (temperature, precipitation, pressure, evaporation, etc.), the temporal resolution of the observations (annual, seasonal, monthly or sub-monthly), the availability of metadata (station's history information) and the monitoring station network density (spatial resolution). A review of different methods for the homogenization of climate series is presented by Peterson *et al.* (1998), and comparisons between procedures are provided by Ducré-Robitaille *et al.* (2003) and Reeves *et al.* (2007). Following the hybrid approach proposed by Wijngaard *et al.* (2003) for the European Climate Assessment & Dataset (ECA&D) project, we did not attempt to remove non-climatic inhomogeneities from the 107 daily precipitation series compiled, but rather provide a qualitative classification of each station's records. Therefore, the results of the homogenization analysis were used to develop an overall classification of the daily series.

The second part of the article investigates the existence of trends and other temporal patterns in extreme precipitation indices, within the period 1955–1999, at 15 monitoring stations located in southern Portugal. This 45-year period was chosen to optimize data availability across the region, taking into consideration the homogenization analysis performed. In all, three of the indices (SDII, R5D and R30) provide information on the 'wetness', whereas the other three [CDD, AII and frequency of dry spells (FDD)] characterize the 'dryness'. The selected indices are appropriate for the purposes of this research, because they might contribute to assess climate dynamics that must be accounted for in impact studies related with water resources management, environmental policies, land use and desertification-related studies for the South of Portugal. The six daily precipitations indices were analysed through regression models and smoothing techniques.

This article is organized in two major parts. The first one (Section 2) addresses the homogenization assessment of the daily precipitation series, and the second part (Section 3) aims to characterize the dynamic temporal evolution of extreme precipitation indices in the 1955–1999 period. Finally, Section 4 states the major conclusions.

## 2. Daily rainfall database and quality control

### 2.1. Study domain and precipitation data

The study domain refers to the South of continental Portugal, and is defined by the Arade, Guadiana, Mira, Ribeiros do Algarve and Sado basins. The daily precipitation series analysed were compiled from the European climate assessment (ECA) dataset and the National System of Water Resources Information (Sistema Nacional de Informação de Recursos Hídricos (SNIRH), managed by the Portuguese Institute for Water) database, and are available through free downloads from the ECA&D project website (<http://eca.knmi.nl>) and the SNIRH website (<http://snirh.inag.pt>), respectively. The analysed precipitation series were downloaded during the first semester of 2004. Despite being outside the study domain, data from Lisbon and Badajoz (Spain) stations were also compiled from the ECA dataset. All stations with at least 30 years with less than 5% of observations missing were selected. Shorter series with at least 10 years lacking a maximum of 5% of data were also chosen, and hence the series with too many gaps were discarded. Using those criteria, 45 long-term and 62 short-term series of daily precipitation were accepted for the homogenization analysis. Even though the beginning and ending of series from the SNIRH database are highly variable, 44 long-term series have a common period of observation of 20 years, located in the 1964/1983 interval. Most of the long-term series (more than 90%) cover the standard normal period 1961/1990, and 33% of them extend back to 1931. Figure 1 shows the study domain and the geographical distribution of stations for which daily time series have been selected. The data are spatially representative of the study domain that covers approximately 25 200 km<sup>2</sup>.

Before being collected for this study, the daily series of the ECA dataset had already been subject to several basic quality-control procedures and statistical homogeneity testing. Because of the sparse density of the ECA station network, absolute tests were applied rather than relative tests, i.e. testing candidate station's series relative to neighbouring stations' series, which are presumed homogeneous. The ECA&D project used historic metadata information to find supporting evidence of changes in observational routines that may have triggered the irregularities detected. The ECA daily series were not adjusted for the inhomogeneities identified. Instead, the results of the different tests were grouped in an overall classification ('useful', 'doubtful' and 'suspect'). The four long-term precipitation series [Beja (666), Lisboa Geofísica




Figure 1. Study area and stations with daily precipitation series. Station dots are scaled with the length of the time series. Red dots: long-term series. Black dots: short-term series. This figure is available in colour online at [www.interscience.wiley.com/ijoc](http://www.interscience.wiley.com/ijoc)

(675), Tavira (681) and Badajoz Talavera (709)] compiled from the ECA dataset for this study were all marked as 'useful', as the four homogeneity tests did not reject the homogeneity hypothesis, at the 1% level (ECA&D project, <http://eca.knmi.nl>; Klein Tank *et al.*, 2002; Wijnngaard *et al.*, 2003).

Some homogeneity testing of the annual precipitation totals of the stations from the SNIRH database has been carried out by Nicolau (1999), for the period 1959/1960–1990/1991. This author performed a double-mass analysis and three absolute homogeneity tests. Nicolau (1999) found no inhomogeneities in the annual precipitation series of the monitoring stations considered here. In summary, the full length of the series from the SNIRH database was not analysed and *objective* relative methods were not performed. Therefore, we assumed that the selected 107 daily precipitation series could contain potential breaks, as recommended by Auer *et al.* (2005), and thus several homogeneity testing procedures were applied to all of them.

## 2.2. Homogeneity assessment methodology

There are a number of tests available for the homogenization of climate series with low temporal resolution (e.g. Peterson *et al.*, 1998). However, well-established statistical methods for the homogeneity testing of sub-monthly precipitation data are lacking (Wijnngaard *et al.*, 2003; Auer *et al.*, 2005). Furthermore, adjusting daily and hourly data is not straightforward, thus the World Meteorological Organization (WMO) makes no recommendations regarding adjusting sub-monthly data (Aguilar *et al.*, 2003). In order to overcome those limitations and taking into consideration the previous quality control-analysis of the selected ECA series, the homogeneity assessment followed the hybrid approach proposed by Wijnngaard *et al.* (2003) for the ECA dataset. Hence, the homogeneity procedures used as the testing variable, the annual wet day count with 1-mm threshold, which is expected to be representative of important characteristics of variation at the daily scale. The results of the different

procedures implemented were then used to develop an overall classification of the daily series.

The homogeneity assessment of the precipitation time series was developed through four major stages (Figure 2). The first one comprises several basic quality control-procedures that aim at the identification of errors and suspicious daily precipitation records, which were flagged using several criteria. The second stage is dedicated to absolute homogeneity testing and comprises the application of six statistical tests to the testing variable, at all locations: the Mann–Kendall test (Mann, 1945; Kendall, 1975), the Wald–Wolfowitz runs test (Wald and Wolfowitz, 1943), the Von Neumann ratio test (Von Neumann, 1941), the Standard normal homogeneity test (SNHT) for a single break (Alexandersson, 1986), the Pettit test (Pettit, 1979) and the Buishand range test (Buishand, 1982). In order to select a subset of series with quality data, the outcomes from the six tests were then grouped together, and a classification was established relying on the number of tests rejecting the homogeneity hypothesis at the 5% significance level. For the long-term series, the criteria were the following: (1) series considered homogeneous by all tests were classified as 'reference'; (2) series for which only one of the six tests rejected the null hypothesis were classified as 'candidate'; (3) series for which two or more absolute tests rejected the homogeneity hypothesis were not analysed further. In the relative testing stage, the selected reference series were also tested through an iterative procedure in which they were seen consecutively as candidates and references. For the short-term series, two criteria were considered: (1) series considered homogeneous by all tests were classified as 'useful'; (2) series for which at least one of the absolute tests rejected the homogeneity hypothesis were classified as 'doubtful'.

The relative testing stage comprises the application of those last three homogeneity tests to long-term composite ratio series (Alexandersson and Moberg, 1997), and the application of a new procedure to the testing variable. This technique is an extension of the Ellipse test,




Figure 2. Schematic representation of the methodology for the homogeneity assessment of the precipitation time series.

described by Allen *et al.* (1998), that takes into account the contemporaneous relationship between several candidate series from the same climatic area by using the residuals from a *seemingly unrelated regression equations* (SUR) model (Zellner, 1962), thus named SUR + Ellipse test (Appendix).

Finally, in the fourth stage, a geostatistical stochastic simulation approach was applied to the testing variable of four candidate stations (Costa *et al.*, 2008). This procedure uses the direct sequential simulation algorithm (Soares, 2001) to determine local probability density functions at candidate stations' locations, by using spatial and temporal neighbouring observations. The results suggest that this procedure allows for the identification of breakpoints near the start and end of a series, and allows for the detection of multiple breaks simultaneously (Costa *et al.*, 2008). All other testing techniques considered were used iteratively by systematically dividing the tested series into smaller segments when a break was detected, and then performing the test on those segments.

### 2.3. Homogeneity assessment results

All statistical tests results used the 5% significance level, and the data analysis was generated through specific programs developed using SAS software macros, SAS/STAT, SAS/ETS and SAS/GRAPH software of the SAS System (registered trademarks of SAS Institute Inc.) for Windows, Version 8.

#### 2.3.1. Basic quality control analysis

Routine quality-control procedures revealed that all precipitation records were non-negative but many series had non-existent dates, which were properly corrected and missing values were assigned to the variable for those days. Several robust location and scale estimates were computed for outlier detection by using all records from the daily time series, and by computing estimates for each year. The upper asymmetric pseudo-standard deviation (Lanzante, 1996) was computed for all 107 daily precipitation series, but it was inconclusive since the median is equal to zero for all series and the third quartile is different from zero for 11 series only, thus the interquartile range is always equal to zero except for those 11 series. The biweight estimates of the mean and standard deviation (Lanzante, 1996) could not be computed because the median-absolute-deviation (MAD) is equal to zero for all daily precipitation series and it appears in the weights denominator of those estimates. Similarly, Feng *et al.* (2004) applied this procedure for temperature data only. Robust alternatives to MAD are the  $S_n$ - and  $Q_n$ -standard deviations (Rousseeuw and Croux, 1993). The  $S_n$ -standard deviation is equal to zero for all daily precipitation series, and the  $Q_n$ -standard deviation is approximately equal to 0.22 mm for all series, indicating that the centres of the distributions have low variability.

The next set of procedures aimed to identify questionable data by flagging the daily precipitation records

with the following classification scheme: (1) 'useful', (2) 'doubtful', (3) 'suspect' and (4) 'erroneous'. The first criterion relied on data outlying pre-fixed thresholds: records greater than the 99th percentile were flagged as (2); records greater than 100 mm were flagged as (3) and all others as (1). Using this criterion, 44% from the whole 107 series under analysis had records flagged as (3), in which 25 of them were long-term series and the other 22 were short-term series. Not surprisingly, the long-term series had an average number of records flagged as (3) approximately equal to 5, and for the short-term ones that average was approximately 3. The total number of records flagged as (3) was equal to 188. The second criterion used was a subjective evaluation of data previously flagged as (3), 'suspect'. If at least two monitoring stations had daily precipitation records greater than 100 mm on the same day, or within a 1-day range, their flag was set to (2), 'doubtful'. As a result, the number of records flagged as (3) dropped to 52 and the number of series to 22 (16 long-term and 6 short-term).

The third criterion relied on graphical analysis. All 107 series were plotted against time, and when a peak in the graph seemed suspicious, even if that value was previously classified as (1) or (2), a closer look was taken by plotting the data against time together with highly correlated stations (Pearson's correlation coefficient greater than 0.70 or highly significant Spearman rank-order correlation coefficient) for the 3-month period centred in the suspicious day. After a subjective analysis of all the graphs (over 500), several records were reclassified. Afterwards, the Portuguese Institute for Water (INAG – Instituto da Água) was contacted in order to clarify if data flagged as (4), 'erroneous' were outliers or a result of extreme weather phenomena. The erroneous values identified were then set to missing. Among the series with records flagged, the most problematic ones are Alcoutim (29M.01) and Picota (30K.02), both from the SNIRH database. It might be advisable to set to missing, the daily records of the years 1954–1959 of Alcoutim, as they were found highly suspicious. The daily precipitation records of December 1972 and December 1973 are precisely the same in Picota, thus it might also be advisable to set them to missing.

The last quality-control procedure was a 'flat line' check (Feng *et al.*, 2004), which identifies data of the same value for at least 3 consecutive days (not applied to zero precipitation data). For those detected records, the first occurrence was flagged as (0) 'useful', and the following records as (1) 'suspect'. All other records were flagged as (0) 'useful'. Almost half (49%) of the long-term series and 19% of the short-term ones were flagged with 'suspect' records using this methodology. The average number of *runs* (blocks of 3 or 4 consecutive days having the same value) per station was equal to two. The flagged precipitation values range from 0.1 to 5 mm and the most common values were 0.1 and 0.2 mm. This seems to indicate that if those flagged values are erroneous they might have been originated by measurement errors (i.e. how precisely very low amounts

of precipitation are measured) rather than by editing errors.

### 2.3.2. Absolute testing

The absolute testing stage comprises the application of six statistical tests to the testing variable at the 107 monitoring stations. Two of the homogeneity tests applied are not distribution free, namely the SNHT and the Buishand range test, and assume that data are independent, identically normally distributed random quantities. Moreover, the remaining non-parametric tests applied also require serially independent data. For those reasons, generalized Durbin–Watson autocorrelation tests and four normality tests were applied to the testing variable series at all stations.

The Durbin–Watson test is a widely used method of testing for autocorrelation. The generalized Durbin–Watson statistics for 1st, 2nd and 3rd order autocorrelation were computed, and conclusions were drawn at the 5% level. The generalized Durbin–Watson tests revealed 1st-order autocorrelation for almost 19% of the series (16 long-term and 4 short-term), and 2nd-order for four series only. None of the testing series had significant 3rd-order autocorrelation. The four normality tests applied were the Shapiro–Wilk, the Kolmogorov–Smirnov, the Cramér–von Mises and the Anderson–Darling tests. For details on the statistical computation of the normality tests refer to SAS Institute (1999, pp. 1397–1401). In view of the results from those four normality tests, over 80% of the testing series (36 long-term and 50 short-term) were considered as Gaussian by all of them. On the other hand, the four tests rejected the normality hypothesis for 7.5% of the series (2 long-term and 6 short-term). Taking into consideration these results, we decided to proceed with the homogeneity tests. Moreover, it is a standard procedure to relax those assumptions for annual data.

Regarding the homogeneity testing results (Table I), approximately 38% of the long-term series were considered appropriate to be selected as reference, and 24% as candidate. Thus, the remaining 38% were excluded from the relative testing analysis. Not surprisingly, approximately 76% of the short-term series were considered homogeneous by the six statistical tests, and thus globally evaluated as 'useful'.

### 2.3.3. Relative testing

The results from the relative testing stage are detailed in Table II. The series from Viana do Alentejo (24I.01) were not tested using the SUR + Ellipse test because it was not possible to determine a common period, without too many gaps, for all the series that would be appropriate to model simultaneously (candidates and their respective references). All the regressors (reference series) parameters of the SUR models are statistically significant. Each SUR model includes at least two candidate stations' data, and some series were tested more than once through different models, depending on the common period of the

Table I. Results from the absolute testing stage and overall classification of the daily series. The Mann–Kendall (MK), Wald–Wolfowitz (WW), Von Neumann (VN), SNHT, Pettit (P) and Buishand (B) tests were applied to the annual number of wet days (threshold 1 mm), and used the 5% significance level.

Station	Code	Period	Tests rejecting the homogeneity hypothesis and break years detected	Classification
Tavira	681	1941–1994	MK; P (1979); SNHT (1978)	Potentially suspect
São Julião	18N.01	1981–1999	Homogeneous	Potentially useful
Alegrete	18N.02	1981–1999	Homogeneous	Potentially useful
Santa Eulália	19N.02	1983–1999	Homogeneous	Potentially useful
Esperança	19N.03	1980–1999	Homogeneous	Potentially useful
Degolados	19O.03	1984–1999	Homogeneous	Potentially useful
Caia (M. Caldeiras)	20O.02	1980–1999	SNHT (1989)	Potentially useful
Vendas Novas	21G.01	1932–1999	VN; SNHT (1943)	Potentially useful
Vila Viçosa	21M.01	1981–2000	MK	Potentially suspect
Alandroal	21M.02	1984–1999	SNHT (1989)	Potentially suspect
Juromenha	21N.01	1932–1999	MK; VN	Potentially useful
Águas de Moura	22E.01	1984–2001	Homogeneous	Potentially useful
Moinhola	22F.03	1973–2000	VN	Potentially useful
Santa Susana	22L.02	1950–1999	MK; VN; B, P and SNHT (1979)	Potentially suspect
Santiago Maior	22M.01	1984–1999	Homogeneous	Potentially useful
Montevil	23F.01	1984–2000	Homogeneous	Potentially useful
Barragem de Pego do Altar	23G.01	1980–2000	Homogeneous	Potentially useful
Reguengos	23L.01	1985–1999	Homogeneous	Potentially useful
Grândola	24F.01	1973–2000	MK; VN; P (1979); SNHT (1982)	Potentially suspect
Barragem do Vale do Gaio	24H.02	1980–2000	VN	Potentially suspect
Barragem de Odivelas	24I.03	1974–2000	VN	Potentially suspect
Alvito	24J.02	1984–2000	Homogeneous	Potentially useful
Cuba	24J.03	1986–1998	Homogeneous	Potentially useful
Portel	24K.01	1984–1999	Homogeneous	Potentially useful
Vidigueira	24K.02	1984–2000	Homogeneous	Potentially useful
Amareleja (D.G.R.N.)	24N.01	1984–2000	Homogeneous	Potentially useful
Ferreira do Alentejo	25I.01	1933–2000	VN; B, P and SNHT (1958)	Potentially suspect
Pedrogão do Alentejo	25L.01	1942–2000	WW; VN; B (1954); P (1954, 1972); SNHT (1946, 1954, 1965)	Potentially suspect
Sobral da Adiça	25N.01	1981–2000	SNHT (1983)	Potentially suspect
Santo Aleixo da Restauração	25O.01	1932–2000	WW; VN; B and P (1958)	Potentially useful
Barrancos	25P.01	1986–1998	Homogeneous	Potentially useful
Barragem de Campilhas	26F.02	1956–1994	VN; P (1979, 1989); SNHT (1979)	Potentially useful
Santa Vitória	26I.01	1984–2000	Homogeneous	Potentially useful
Aljustrel	26I.03	1936–2000	WW; MK; VN; B, P and SNHT (1972)	Potentially useful
Albernoa	26J.04	1984–2000	Homogeneous	Potentially useful
Salvada	26K.01	1984–2000	Homogeneous	Potentially useful
Serpa	26L.01	1985–2000	Homogeneous	Potentially useful
Santa Iria	26L.02	1980–2000	Homogeneous	Potentially useful
Garvão (Montinho)	27G.02	1980–1994	SNHT (1983)	Potentially suspect
Barragem do Monte da Rocha	27H.02	1980–1994	Homogeneous	Potentially useful
Castro Verde	27I.01	1932–2000	MK; P (1979)	Potentially useful
São Marcos da Ataboeira	27J.01	1984–2000	Homogeneous	Potentially useful
Corte Pequena	27J.02	1986–1996	Homogeneous	Potentially useful
Vale de Camelos	27J.03	1990–2000	Homogeneous	Potentially useful
Algodôr	27K.01	1987–1999	Homogeneous	Potentially useful
Corte da Velha	27K.02	1980–2000	Homogeneous	Potentially useful
Barragem de Mira	28G.01	1970–1993	VN	Potentially suspect
Santana da Serra	28H.03	1936–2000	MK; P (1979)	Potentially useful
Almodôvar	28I.01	1984–2000	Homogeneous	Potentially useful
Alcaria Longa	28J.01	1986–1999	Homogeneous	Potentially useful
Santa Barbara de Padrões	28J.03	1980–2000	Homogeneous	Potentially useful
São João dos Caldeireiros	28K.01	1984–2000	Homogeneous	Potentially useful
Álamo	28K.02	1981–2000	Homogeneous	Potentially useful
Mértola	28L.01	1984–2000	Homogeneous	Potentially useful
Cimalhas	29F.01	1981–1999	SNHT (1982)	Potentially suspect

Table I. (Continued).

Station	Code	Period	Tests rejecting the homogeneity hypothesis and break years detected	Classification
Foz do Farelo	29F.02	1981–1999	SNHT (1983)	Potentially suspect
São Barnabé	29I.01	1965–2000	VN; P (1972)	Potentially suspect
Santa Clara-a-Nova	29I.02	1981–1999	Homogeneous	Potentially useful
Guedelhas	29J.05	1980–1999	Homogeneous	Potentially useful
Martim Longo	29K.01	1985–2000	Homogeneous	Potentially useful
Malfrades	29K.03	1981–1999	Homogeneous	Potentially useful
Penedos	29K.04	1981–2000	MK; SNHT (1995)	Potentially useful
Pereiro	29L.01	1958–1999	VN; B (1983); SNHT (1995)	Potentially suspect
Monte dos Fortes	29L.03	1984–1999	Homogeneous	Potentially useful
Alcoutim	29M.01	1939–1999	WW; VN; B, P and SNHT (1959)	Potentially suspect
Marmelete	30E.02	1984–1999	Homogeneous	Potentially useful
Monchique	30F.01	1933–1998	WW; VN	Potentially useful
São Bartolomeu de Messines	30H.03	1991–1998	Homogeneous	Potentially useful
Paderne	30H.05	1984–1999	MK	Potentially suspect
Sobreira	30I.02	1943–1999	WW; VN; B (1954); SNHT (1949)	Potentially useful
Mercador	30K.01	1984–1999	Homogeneous	Potentially useful
Faz-Fato	30L.03	1956–1999	VN; B and P (1986)	Potentially suspect
Lagos	31E.01	1956–1999	MK; B and SNHT (1972); P (1979)	Potentially suspect
Porches	31G.02	1980–1998	Homogeneous	Potentially useful
Algoz	31H.02	1981–1996	Homogeneous	Potentially useful
São Brás de Alportel	31J.01	1985–2000	Homogeneous	Potentially useful
Estoi	31J.04	1984–1999	Homogeneous	Potentially useful
Santa Catarina (Tavira)	31K.01	1984–1999	Homogeneous	Potentially useful
Quelfes	31K.02	1982–1998	Homogeneous	Potentially useful

Table II. Results from the relative testing stage and overall classification of the daily series. The Buishand, Pettit, and SNHT tests were applied to composite (ratio) reference series. The SUR + Ellipse test and the geostatistical simulation approach (Costa *et al.*, 2008) were applied to the annual number of wet days (threshold 1 mm). All tests used the 5% significance level.

Station	Code	Period	Relative tests rejecting the homogeneity hypothesis and break years detected	Classification
Beja	666	1951–1999	Geostatistical simulation approach: 1991	Suspect
Lisboa Geofísica	675	1941–1999	Homogeneous	Useful
Badajoz Talavera (Spain)	709	1955–2000	Pettit test: 1975	Suspect
Arronches	19N.01	1932–1999	SNHT: 1954 SUR + Ellipse test: 1988	Suspect
Barragem do Caia	19O.02	1965–2000	Homogeneous	Useful
Azaruja	21K.01	1944–1982	Homogeneous	Useful
Santiago do Escoural	22H.02	1932–1999	SUR + Ellipse test: 1960 Buishand and Pettit tests: 1988 SNHT: 1989 Geostatistical simulation approach: 1987, 1988, 1996	Suspect
Redondo	22L.01	1945–1982	Buishand, Pettit, SNHT, SUR + Ellipse tests: 1963	Suspect
Comporta	23E.01	1934–2000	Pettit test, SNHT: 1986	Useful
Alcáçovas	23I.01	1932–2000	Buishand, Pettit, SNHT, SUR + Ellipse tests: 1960	Suspect
São Manços	23K.01	1943–2000	SNHT and SUR + Ellipse test: 1950	Suspect
Viana do Alentejo	24I.01	1934–2000	Homogeneous	Useful
Azinheira Barros	25G.01	1951–2000	Homogeneous	Useful
Barragem do Roxo	26I.02	1959–2000	Homogeneous	Useful
Herdade de Valada	26M.01	1969–2000	SNHT: 1995	Suspect
Relíquias	27G.01	1932–2000	Buishand and Pettit tests: 1969	Suspect
Panóias	27H.01	1956–1994	Homogeneous	Useful
Odemira	28F.01	1932–1994	SUR + Ellipse test: 1952	Useful
Aldeia de Palheiros	28H.01	1932–1996	Homogeneous	Useful
Sabóia	29G.01	1932–1994	Buishand test: 1949 Buishand, Pettit, SUR + Ellipse tests: 1984 SNHT: 1985	Suspect

Table II. (Continued).

Station	Code	Period	Relative tests rejecting the homogeneity hypothesis and break years detected	Classification
Aljezur	30E.01	1932–1999	Buishand, Pettit, SNHT, SUR + Ellipse tests: 1968	Suspect
Barragem da Bravura	30E.03	1956–2000	Homogeneous	Useful
Alferce	30G.01	1959–1999	Buishand, Pettit, SUR + Ellipse tests: 1984 Geostatistical simulation approach: 1983	Suspect
Santa Margarida	30H.04	1965–1999	Buishand and Pettit tests, SNHT: 1978	Suspect
Barranco do Velho	30J.01	1956–1999	Buishand and Pettit tests: 1976 SNHT: 1975, 1996	Suspect
Catraia	30J.02	1932–1973	Homogeneous	Useful
Picota	30K.02	1957–1999	Buishand test: 1988	Useful
Alcaria (Castro Marim)	30L.04	1947–1999	Homogeneous	Useful




Figure 3. SUR + Ellipse test results for the annual number of wet days (threshold 1 mm) at Aljezur (30E.01) station (left graph: testing period is 1941/99; right graph: testing period is 1956/97).

series included in each model. Consequently, depending on the testing period, different models sometimes provided different results for a specific candidate series. This problem can be minimized by testing the candidate series through a different model whenever a peak in the graph from the Ellipse test seems suspicious (Figure 3, left graph), or by using a combination of statistical tests. In fact, Wijngaard *et al.* (2003) state that, generally, a combination of statistical methods and methods relying on metadata information is considered to be most effective to track down inhomogeneities. Nevertheless, that problem might also happen with other testing methods, although it is harder to detect because the tests are usually applied only once. For example, applying the SNHT to the testing variable of Vendas Novas (21G.01) station for the period 1932/1992 concludes the series as homogeneous, but testing the period 1938/1999 identifies a break in 1943.

The relative magnitudes of the breaks detected are given by the ratio between the average annual wet day count before and after two consecutive breaks. The magnitudes of the breaks detected by the SUR + Ellipse test, but not identified by the other methods, range from  $-7.6\%$  to  $6.88\%$ . Conversely, the magnitudes of the breaks detected by at least one of the other three

tests, but not identified by the SUR + Ellipse test range from  $-14.09$  to  $10.78\%$ . Hence, there is no apparent connection between the potential breaks magnitudes and the ability of the SUR + Ellipse test to identify them.

Only 4 of the 11 series previously classified as candidates were considered as homogeneous by all the relative tests. Considering the 17 series previously classified as references, 8 of them were considered as homogeneous by all the relative tests. The breaks detected are mainly located between 1949 and 1954, and around 1986. Therefore, there is an apparent trend towards less breaks in recent times, in contrast to that reported by other homogenization studies (Tuomenvirta, 2001; Wijngaard *et al.*, 2003; Auer *et al.*, 2005). The station selection was on the basis of the absolute testing results, thus that apparent trend may not be true if all the 107 stations' series were tested through the relative approach.

#### 2.3.4. Overall classification and discussion

An overall classification of the daily precipitation series was established using four classes (Table III): 'useful', 'potentially useful', 'potentially suspect' and 'suspect'. A series was classified as 'useful' when all relative approaches (the four relative statistical tests and the stochastic approach) considered it as homogeneous.



Table III. Criteria used to establish the overall classification of the daily series.

Classification	Criteria
<i>Useful</i>	All relative approaches considered the series as homogeneous. Relative break(s) detected might be explained by several months without records.
<i>Potentially useful</i>	Short-term series previously classified as 'useful' (the six absolute tests considered the series as homogeneous). Absolute break(s) detected might be explained by several months without records.
<i>Potentially suspect</i>	Absolute break(s) detected could not be explained by non-climatic factors.
<i>Suspect</i>	Relative break(s) detected could not be explained by non-climatic factors.

Whenever the daily series had several months without records near a break year, identified by some relative testing procedure, the series was also classified as 'useful', because it is conceivable that the inhomogeneous records were set to missing in the SNIRH database, and the tests rejections were due to them. A series was classified as 'suspect' when at least one of the relative approaches considered it as inhomogeneous and the break(s) detected could not be explained by non-climatic factors.

Considering the series analysed through absolute testing only (both short and long-term), it is difficult to determine if changes or lack of changes result from non-climatic or climatic influences (Peterson *et al.*, 1998), since it was not possible to find historic metadata support. Therefore, the intermediate classes, 'potentially useful' and 'potentially suspect', were established. Furthermore, as the short-term series were only analysed through absolute testing, those series were classified as 'potentially useful' if the six absolute tests considered the series as homogeneous. Relative approaches that use data from reference stations are usually preferred because they aim to isolate the effects of station irregularities and to account for regional climate changes. In fact, the results show that the relative tests identified inhomogeneities in a number of stations that were previously considered as homogeneous by the six absolute tests, and vice-versa. Although desirable, a relative approach for the homogeneity assessment of all series could not be used since it was out of the scope of this research.

Following those criteria, approximately 13% of the 107 series were classified as 'useful', 55% were classified as 'potentially useful', 19% were classified as 'potentially suspect' and 13% as 'suspect' (Tables I and II). Although defined with different criteria, the qualitative interpretation of the overall classes is similar to the one given for the categories defined for the ECA series (Wijngaard *et al.*, 2003). However, it is important to point out that we used the 5% significance level in all statistical tests, whereas those authors used the 1% level. Therefore, our classification is more conservative in the sense that we allowed for the rejection of the homogeneity hypothesis at stations that are considered homogeneous at the 1% significance level. The series classified as 'useful' seem to be sufficiently homogeneous for trend analysis and variability analysis. The series classified as 'potentially

'useful' and 'potentially suspect' should be used cautiously, from the perspective of the existence of possible inhomogeneities, as the homogeneity analysis performed might be considered inconclusive – even though all series were considered homogeneous by previous studies. The series classified as 'suspect' should be excluded from trend analysis and variability analysis, as there is strong evidence of inhomogeneities present.

### 3. Trends in indices of daily extreme precipitation

Numerous extreme precipitation indices are described and analysed in the literature (Peterson *et al.*, 2001; Frich *et al.*, 2002; Kiktev *et al.*, 2003; Klein Tank and Können, 2003; Haylock and Goodess, 2004; Kostopoulou and Jones, 2005; Moberg and Jones, 2005). There are two main categories of extremes indices: those based on either absolute thresholds or percentiles. The first category refers to counts of days crossing a specified absolute value (e.g. the number of days per year with daily precipitation exceeding 30 mm). The second category of indices is on the basis of statistical quantities such as percentiles, so the tails of the statistical distribution are examined and days exceeding (not exceeding) a given high (low) percentile are counted. Indices based on percentile thresholds have a clear advantage for climate-change detection studies as they compare the changes in the same parts of the precipitation distributions and thus can be used in studies of wide regions (Haylock and Nicholls, 2000; Klein Tank and Können, 2003). On the other hand, indices based on the count of days crossing certain fixed thresholds are beneficial for impact studies as they can be related with extreme events that affect human society and the natural environment (Klein Tank and Können, 2003).

The later set of indices, and indices describing events with short return periods (moderate climate extremes), are suitable for the purposes of this research since they might contribute to assess climate dynamics at the local scale that contribute for land degradation and desertification prone areas of the South of Portugal. Accordingly, we selected four extreme precipitation indices recommended by the joint CCI/CLIVAR/JCOMM Expert Team on Climate Change-Detection and Indices (ETCCDI, <http://www.clivar.org/organization/etccdi/etccdi.php>; Peterson *et al.*, 2001; Frich *et al.*, 2002), and developed

two other indices describing dry conditions. The following sections explain the criteria for station selection from the developed daily rainfall database, the indices rationale and definitions, the trend analysis methodology and, in the last section, the results are summarized and discussed.

### 3.1. Analysis period and data selection

From the set of 107 stations compiled for homogeneity assessment, one station's data (29G.01 Sabóia) were excluded from the analysis because multiple breakpoints were identified and the homogeneous periods were too short and unreliable; the Badajoz Talavera (709) station, in Spain, was also excluded. The daily rainfall database for the South of Portugal comprises records in the period 1931/2000, but the beginning and ending of each series are highly variable. The selection of stations with quality data for a long common period was developed through several stages.

First, the extreme precipitation indices were computed for the set of 105 stations, regardless of their overall homogeneity classification. Nevertheless, only the longest homogeneous period was used to build the indices for the series classified as 'suspect'. The extreme precipitation indices are sensitive to the number of missing days, thus the daily records of the selected stations should be as complete as possible. Consequently, for each station, the indices for a specific year were set to missing if there were more than 16% of the days missing for that year (Haylock and Goodess, 2004).

Next, a first set of stations was selected for trend analysis by including all the series classified as 'potentially useful', 'useful' and the longest homogeneous period of the series classified as 'suspect'. In this set, the number of stations with at least 30 years of overlapping observations was very small. Hence, the next stage aimed to select stations classified as 'potentially suspect' with break years near the beginning of the series (identified through absolute testing), so that their longest homogeneous period could also be considered. This allowed us to determine the analysis period 1955/1999, which is the longest common period for the final set of 15 series. The regional analysis of anomalies used four additional series with homogeneous records within 1940/2000 (Figure 4). All stations selected have less than 12% of the days missing in each year, and the data for most stations do not have any missing records.

### 3.2. Precipitation indices

In the present study only annually specified indices are considered. Their definitions are listed in Table IV. The SDII is a simple daily intensity index defined as the average precipitation per wet day and a wet day is defined as a day with at least 1 mm of precipitation. The R5D index is defined as the highest consecutive 5-day precipitation total and can be considered a flood indicator, since it provides a measure of short-term precipitation intensity. The R30 index characterizes the frequency




Figure 4. Stations selected for trend analysis. Dots: stations with nearly complete records in 1955–1999. Triangles: additional stations with data within 1940–2000 used to build the regional-average anomaly time series.

of extremely heavy precipitation events and is defined as the number of days with daily precipitation totals above or equal to 30 mm. This threshold fits the extreme events regime of the study area as the 30-mm value approximately corresponds to the 95% regional-average percentile of the 1961/90 climate normal. The CDD index corresponds to the maximum number of consecutive dry days, and therefore characterizes the length of the greatest dry spell.

Not only drought but also moderate dry conditions have significant impacts in terms of crop losses, water supply shortages, land degradation and desertification in the South of Portugal. To better understand the pluviometric regime of this region, and the frequency and magnitude of dryness in particular, we developed two indices describing dry events (FDD and AII).

The FDD index is defined as the number of dry spells. For return periods of 2 years, expected dry-lengths vary from 60 to 80 days in the study region (Lana *et al.*, 2008). The selected indices refer to precipitation events with return periods typically of less than 1 year, providing relevant information to impact studies. In the FDD definition, a dry spell is a consecutive period with at least 8 dry days. The average length of dry spells in a year ranges from 8 to 12 days in the study region (Lana *et al.*, 2008). Therefore, increasing (decreasing) trends of FDD are indicators of a change in the mean frequency of dry events, rather than in the frequency of extremely dry situations, although a change in the extreme values of the distribution obviously implies a change in the mean.

Regarding the SDII, CDD and FDD indices, a wet day is defined as a day with at least 1 mm of precipitation ( $R \geq 1$  mm), thus a dry day has less than 1 mm of precipitation ( $R < 1$  mm). Ceballos *et al.* (2004) state that rainfall amounts below this threshold are not absorbed by soils and are evaporated off directly. In fact, Moberg and

Table IV. Acronyms and definitions of the six indices for moderate precipitation extremes.

Acronyms	Explanation	Units
SDII	Ratio between the total rain on wet days and the number of wet days ( $R \geq 1$ mm).	mm
R5D	Highest consecutive 5-day precipitation total.	mm
R30	Number of days with daily precipitation totals above or equal to 30 mm.	Days
CDD	Maximum number of consecutive dry days ( $R < 1$ mm).	Days
FDD	Number of dry spells (consecutive period with at least 8 dry days, $R < 1$ mm).	Freq.
AII	Ratio between the total rain on dry days and the number of dry days ( $R < 10$ mm).	mm

Jones (2005) agree that, with this definition, a dry day is allowed to have a small amount of precipitation, but generally small enough as the ground will not recover after a long period of dryness. Moreover, thresholds lower than 1 mm can introduce trends in the number of wet days, associated with measurement errors introduced by the observers (Haylock and Nicholls, 2000; Haylock and Goodess, 2004) or by instrument inaccuracies. In fact, taking into consideration the ‘flat line’ check results, it is prudent to adopt such a threshold for dry days ( $R < 1$  mm) because it allows for minimizing any inaccuracy associated with measurement errors.

In the definition of the AII index, we used the 10-mm threshold to indicate a dry day (Ceballos *et al.*, 2004; Lana *et al.*, 2008). Let  $RL10t$  be the total rain on days with precipitation amount below 10 mm ( $R < 10$  mm), and let  $RL10$  be the number of days with  $R < 10$  mm. Similar to the SDII, the AII index is defined by  $RL10t/RL10$  and can be interpreted as a simple aridity index, because it is a numerical indicator of the degree of dryness of the climate at a given location. Below the 10-mm threshold, the rainfall has a small effect on the soil water-content, since the rainfall evaporates very quickly and hardly drains into the upper soil layer (Ceballos *et al.*, 2004). Increasing (decreasing) trends of AII are indicators of change in the normal moisture availability, which is a sensitive issue for desertification susceptible regions.

### 3.3. Trend estimation and diagnosis methods

The six daily precipitations indices were subject to a number of diagnosis tests, at each station’s location, in order to verify the existence of autocorrelation and heteroscedasticity of the regression errors. Depending on the tests’ results, the trend estimation was performed using three different regression models. The indices are expressed as annual values  $Y_t$ ,  $t = 1, \dots, T$  with the subscript  $t$  referring to the year (also denoted by  $X_t$ ), and  $T$  is the length of the period covered by the station’s series. Engle’s Lagrange multiplier test for heteroscedasticity (Engle, 1982) allows to test if the regression errors variance has the form  $V(\varepsilon_t) = \sigma_t^2 = \alpha_1 + \alpha_2 X_t$ , where  $\varepsilon_t$  is the disturbance term (error) of the ordinary least squares (OLS) regression;  $\alpha_1$  and  $\alpha_2$  are constant parameters. Whenever the null hypothesis of homoscedasticity was rejected, the following *heteroscedastic linear model* was fitted:

$$Y_t = \beta_1 + \beta_2 X_t + \varepsilon_t, \quad \varepsilon_t \sim N(0, \sigma_t^2)$$

$$\sigma_t^2 = \sigma^2(\alpha_1 + \alpha_2 X_t) + \eta_t, \quad \eta_t \sim N(0, \sigma_\eta^2) \quad (1)$$

where each error term  $\eta_t$  is normally and independently distributed with mean 0 and constant variance  $\sigma_\eta^2$ . The presence of autocorrelation was investigated using the Durbin–Watson test. Whenever autocorrelation correction was needed, the *autoregressive error model* was fitted:

$$Y_t = \beta_1 + \beta_2 X_t + \varepsilon_t, \\ \varepsilon_t = \rho \varepsilon_{t-1} + \eta_t, \quad \eta_t \sim N(0, \sigma^2) \quad (2)$$

where  $\rho$  is the autoregressive error model parameter. Whenever the homoscedasticity hypothesis and the independent errors assumption were not rejected, the slope of the trend was estimated by OLS. Many stations’ wetness indices had significant non-gaussian residuals (tested through the Shapiro–Wilk test), and other forms of OLS assumptions violations were not investigated (e.g. other forms of heteroscedasticity). Therefore, for all models fitted, the trend significance was assessed through the Mann–Kendall test.

The existence of significant trends in anomaly time series was also investigated using the described methodology. In each year, the anomalies of the indices time series were calculated from the base period of 1961–1990 by standardizing the individual station’s series using the climatological average and standard deviation of the base period. The regional-average anomaly series were computed using the full set of 19 stations’ series (Figure 4) and the analysis period was set to 1940–2000. As all stations do not contain complete data in this period, the regional anomaly of a year was obtained by weighting the anomalies according to the number of stations available for that year (Frich *et al.*, 2002).

Any cyclical components in the variation of time series make it difficult to see the underlying trend. Aiming to improve our understanding of the indices time series, smoothing techniques were used to reduce random fluctuations and provide a clearer view of their underlying behaviour. Moving windows with a time span of 5 and 10 years were used to compute moving average series of the extreme precipitation indices. For the sake of simplicity, these series are denominated the temporal average of extremes (TAE) series. Moving 5 and 10 years, standard deviation statistics of the indices were also computed in order to analyse the temporal evolution

of their variability. These series are denominated the temporal variability of extremes (TVE) series. The TAE and TVE series were calculated for each station and were then averaged over the 15 stations to obtain the regional-average TAE and TVE series for the period 1955/1999.

### 3.4. Results and discussion

A regional correlation analysis, averaging the Spearman rank-order correlation coefficients of the six indices over the 15 stations, revealed that the dryness indices (CDD, FDD, AII) might provide information that is essentially different from the three wetness indices (SDII, R5D, R30), because they are uncorrelated with any of them. Nevertheless, the correlations between AII and the three wetness indices have positive signs, whereas the other two dryness indices show negative signs when correlated with the wetness indices. Interestingly, the correlation between AII and SDII is extremely weak. The correlation between CDD and FDD is negative but weak, which might indicate that an increase (decrease) in the length of the greatest dry spell will not necessarily entail a significant decrease (increase) in the mean frequency of dry events. The three wetness indices are moderately positively correlated with each other.

The results of the Shapiro–Wilk test indicate that most series of the R5D and R30 indices (87% and 93% of the stations, respectively), and 33% of the SDII series, cannot be considered as Gaussian in the period 1955/1999, at the 5% significance level. On the other hand, the normality hypothesis was not rejected for most series of the dryness indices (80% of the CDD series, and 93% of the FDD and AII series).

#### 3.4.1. Trends in extreme indices

Trend estimation results are presented at the 5% and 10% significance levels (Tables V and VI). According to the correlation analysis results, the CDD and FDD indices have oppositely signed trends for most of the stations, although not statistically significant. Hence, these indices do not reflect significant changes neither in the length of dry spells, nor the frequency of dry events. On the other hand, the AII index reflects increases in the magnitude of dryness. A negative station trend in the AII index dominates in the 1955/1999 period, which implies a significant increase of aridity over most of the study region.

The SDII monitors precipitation intensity on wet days and presents significant increasing trends in several stations, but without spatial consistency. A few of them also have significant increasing trends in the maximum 5-day precipitation totals (R5D), but this tendency is not significant for the majority of stations in the R30 index which characterizes the frequency of extremely heavy precipitation events. Accordingly, the extreme indices characterizing wet conditions do not show a clear pattern of significant trends, but rather exhibit different trend signals at the local scale.

These results agree, in general, with those of other studies on regional changes of precipitation for Iberia and Europe (e.g. Klein Tank and Können, 2003; Haylock and Goodess, 2004; Rodrigo and Trigo, 2007), even though the adopted indices and study period do not always coincide and the network of Portuguese stations used in previous studies is coarser. The results from Kostopoulou and Jones (2005), for the period 1958/2000, show regional contrasts over the Eastern Mediterranean in precipitation indices, and significant positive trends were revealed for CDD in many southern stations. Klein Tank and Können (2003) found no significant trends (5% level) in the annual precipitation indices calculated for the southern region of Portugal within the 1946/1999 period. Similarly, the annual indices of precipitation extremes available at the ECA&D project website (<http://eca.knmi.nl>, retrieved 17 March 2008) do not reveal significant trends within the 1946/2006 period, but the scarce number of stations available in southern European regions, especially in Portugal, makes it difficult to assess local contrasts. Nevertheless, the seasonal analysis reveals a few significant (5% level) contrasting trends in the South of Portugal within the 1946/2006 period (<http://eca.knmi.nl>, retrieved 17 March 2008). For example, the frequency of very wet days (R95p) has negative trends at Beja and Tavira in spring; the frequency of extremely wet days (R99p) has a negative trend in Tavira in spring, whereas it has a positive trend in summer (June–August) at Beja. Haylock and Goodess (2004) analysed trends in several extreme precipitation indices over Europe for the winter months (December–February) of the period 1958/2000. In the northwest of the Iberian Peninsula, the results showed a small decrease in CDD, while the rest of the peninsula has seen large increases in this index, except in southern Portugal where the CDD trend magnitudes are small. In contrast, the frequency of very heavy precipitation days (R90p index) showed a decrease over most of the peninsula (including the northwest), but a slight increase in the southeast and southern Portugal. Rodrigo and Trigo (2007) investigated annual and seasonal trends in five precipitation variables using data from 22 stations scattered across the Iberian Peninsula, aiming to analyse the behaviour of daily rainfall in the period 1951–2002. Among those stations, five of them are located in southern Portugal (Lisboa, Grândola, Serpa, Relíquias and Monforte). Only a few significant trends (5% level) were found for these five stations, most of them at Relíquias, and the variables characterizing extreme events do not show a clear pattern at the local scale. For example, Rodrigo and Trigo (2007) showed that the 95th percentile and the percentage of rain falling on days with rainfall above the 95th percentile have negative trends at Relíquias in all seasons and for yearly values, but Serpa has increasing trends for yearly values. Moreover, these variables have significant decreasing trends at Grândola in spring and summer.

The trend results of the anomaly time series (Table VI) are exactly the same as the precipitation indices results (Table V) as far as the trend signals are concerned, but

Table V. Trends in precipitation indices estimated with the *OLS model* (O), the *Autoregressive error model* (A) and with the *Heteroscedastic linear model* (H), for the period 1955/99. Significance of trends assessed using the Mann–Kendall test: values in bold face are significant at <5% level (marked with \*\*) and <10% level (marked with \*).

Station	Code	CDD	FDD	AII	SDII	R5D	R30
Comporta	23E.01	-0.0094	0.0066	(O)	<b>0.0394**</b>	(A)	<b>0.8518**</b> (O) <b>0.0566*</b> (H)
São Manços	23K.01	0.0558	-0.0086	(A)	0.0156	(O)	<b>0.5813*</b> (A) 0.0040 (O)
Azinhreira Barros	25G.01	-0.1734	0.0154	(O)	0.0066	(H)	0.0194 (O)
Ferreira do Alentejo	25I.01	0.1489	-0.0067	(O)	<b>0.0250**</b>	(A)	<b>0.3814*</b> (H) -0.0130 (O)
Pedrogão do Alentejo	25L.01	0.2258	-0.0061	(O)	-0.0117	(A)	0.1358 (H) -0.0093 (A)
Santo Aleixo da Restauração	25O.01	-0.0232	0.0202	(O)	-0.0106	(H)	-0.0144 (A)
Aljustrel	26I.03	<b>0.5369**</b>	-0.0121	(H)	-0.0252	(A)	-0.0066 (A)
Castro Verde	27I.01	-0.2645	<b>0.0497**</b>	(H)	0.0056	(O)	-0.0030 (O)
Aldeia de Palheiros	28H.01	<b>-0.0944**</b>	0.0130	(O)	<b>0.0022**</b>	(H)	-0.0099** (H)
Santana da Serra	28H.03	<b>0.6785**</b>	-0.0228	(O)	<b>0.0636**</b>	(H)	0.0304 (H)
Barragem da Bravura	30E.03	-0.1025	-0.0048	(H)	<b>0.0306*</b>	(O)	<b>0.0617**</b> (H)
Sobreira	30I.02	-0.1640	0.0168	(O)	-0.0021	(O)	<b>0.7732*</b> (A) 0.0377 (H)
Picota	30K.02	-0.3487	0.0141	(A)	<b>0.0588**</b>	(H)	-0.0183 (O)
Alcária (Castro Marim)	30L.04	0.0925	-0.0194	(O)	-0.0238	(O)	-0.0102 (O)
Lisboa Geofísica	675	-0.0939	-0.0080	(H)	-0.0124	(O)	-0.0014 (O)
					-0.0038	(O)	0.0022 (H)

Table VI. Trends in anomaly time series of precipitation indices estimated with the *OLS model* (O), the *Autoregressive error model* (A) and with the *Heteroscedastic linear model* (H), for the period 1955/99. Significance of trends assessed using the Mann–Kendall test: values in bold face are significant at <5% level (marked with \*\*) and <10% level (marked with \*).

Station	Code	CDD	FDD	AII	SDII	R5D	R30
Comporta	23E.01	-0.0003	0.0034	(O)	<b>0.0267**</b>	(A)	<b>0.0263**</b> (O) <b>0.0301*</b> (H)
São Manços	23K.01	0.0021	-0.0049	(A)	-0.0213**	(O)	<b>0.0278*</b> (A) 0.0026 (O)
Azinhreira Barros	25G.01	-0.0083	0.0078	(O)	0.0081	(O)	0.0113 (O)
Ferreira do Alentejo	25I.01	0.0067	-0.0033	(O)	<b>-0.0234*</b>	(H)	<b>0.0243*</b> (H) -0.0113 (O)
Pedrogão do Alentejo	25L.01	0.0083	-0.0031	(O)	<b>-0.0283**</b>	(A)	0.0057 (H) -0.0045 (A)
Santo Aleixo da Restauração	25O.01	-0.0009	0.0095	(O)	-0.0299**	(O)	0.0003 (H) -0.0081 (A)
Aljustrel	26I.03	<b>0.0235**</b>	-0.0044	(H)	-0.0250**	(H)	-0.0196 (O) <b>-0.0041**</b> (A)
Castro Verde	27I.01	-0.0126	<b>0.0273**</b>	(H)	0.0045	(O)	0.0090 (H) <b>-0.0016**</b> (O)
Aldeia de Palheiros	28H.01	-0.0038	0.0060	(O)	<b>-0.0234**</b>	(O)	0.0083 (O) <b>-0.0051**</b> (H)
Santana da Serra	28H.03	<b>0.0259**</b>	-0.0123	(O)	<b>0.0377**</b>	(H)	<b>0.0254*</b> (H) 0.0136 (H)
Barragem da Bravura	30E.03	-0.0037	-0.0024	(H)	-0.0175*	(O)	0.0198 (O) 0.0182 (H)
Sobreira	30I.02	-0.0059	<b>0.0075*</b>	(O)	<b>0.0254**</b>	(H)	<b>0.0224*</b> (A) 0.0104 (H)
Picota	30K.02	<b>-0.0164*</b>	0.0103	(A)	<b>-0.0085*</b>	(O)	-0.0005 (H) -0.0030 (O)
Alcária (Castro Marim)	30L.04	0.0034	-0.0098	(O)	-0.0039	(O)	-0.0006 (O)
Lisboa Geofísica	675	-0.0010	-0.0035	(H)	-0.0166	(O)	0.0055 (H) <b>0.0009*</b> (H)




Figure 5. Differences in the average extreme indices' values between 1940 and 2000 from the average 1961/90 value of weighted regional stations. The trend of the AII annual anomalies series (c) is significant at the 6% level. This figure is available in colour online at [www.interscience.wiley.com/ijoc](http://www.interscience.wiley.com/ijoc)

the trends significance is different for a few stations and indices. Coherent spatial patterns of statistically significant changes emerge in the magnitude of dryness (AII), while the remaining anomaly time series show a lack of spatial consistency. The remaining indicators show mixed patterns of change but significant increases have occurred in the extreme amount derived from short-term precipitation intensity (R5D) in five stations, while for other three stations significant decreases in the number of heavy rainfall events (R30) have occurred. The existence of significant trends in the regional-average anomaly time series was also investigated (Figure 5) using 19 stations' data for the period 1940–2000. As discussed before, absence of spatial consistency and/or significant trends characterizes the majority of the precipitation indices calculated, except for the AII index. Therefore, not surprisingly, this was the only index with a significant decreasing trend (the  $p$ -value of the Mann–Kendall test is equal to 0.06) in the regional-average anomaly time series (Figure 5(c)), indicating an increase of dryness over the study region.

The low spatial coherence of the trends found in the anomaly time series is consistent with the mixed pattern of positive and negative changes shown in the maps of the globe and it is especially noticeable in southern European regions (Frich *et al.*, 2002). For example, the SDII has increased over many parts of Europe, southern Africa, USA and parts of Australia, although the patterns of change are also variable in these parts of the world showing some relatively nearby stations with opposite signs of change (Frich *et al.*, 2002). Absence of spatial consistency also characterizes the majority of the anomaly time series of the precipitation indices for the Eastern Mediterranean (Kostopoulou and Jones, 2005).

### 3.4.2. Dynamic temporal evolution of extremes

Moving window statistics (mean and standard deviation) with a time span of 5 and 10 years were computed for each station, and then averaged over the 15 stations to obtain a regional-average. This is a very useful approach because non-linear trends in precipitation extremes can






Figure 6. Ordinary least squares fitting (OLS, thick line) and weighted local polynomial fitting (LOWESS smoothing, thin line) for each regional-average TAE series (moving average of the extreme index using a time span of 10 years), for the period 1955/99. This figure is available in colour online at [www.interscience.wiley.com/ijoc](http://www.interscience.wiley.com/ijoc)

be revealed, and periods with distinct climatic variability can be identified. The results obtained using windows with a time span of 5 years are identical to the ones obtained with a time span of 10 years, but a little noisier. For this reason, and for the sake of simplicity, only the latter are presented. The moving average series of the extreme precipitation indices was named TAE series. The temporal dynamics underlying these series were captured through *weighted local polynomial models* (LOWESS smoother proposed by Cleveland, 1979) fitted with a time span of  $k$  years determined by generalized cross-validation. In order to point out any non-linear trends underlying the TAE series, simple linear regression models, estimated by OLS, were also fitted.

For all of the precipitation indices considered, the results show non-linear trends in the TAE series within the 1955/1999 period at the large majority of stations.

These results might explain why the regression models fitted to the indices time series could not significantly capture the trend signal for the majority of the stations' indices. Figure 6 shows the results of the regional-average TAE series of the six indices. The non-linear trends and cyclic patterns of the individual stations' TAE series (not shown) are identical to the ones illustrated in Figure 6 (any exceptions are referred in the text), but much more sharpen.

As expected from the previous results, the TAE series of All clearly reflect a strong increase in the magnitude of dryness during the period 1955/1999. The maximum length of dry spells, characterized by the TAE series of CDD, has a decreasing trend until the middle of the 1980s and then suddenly increases, keeping a positive trend through the last decade of the twentieth century. One exception occurs at Pedrogão do Alentejo (25L.01) where

the TAE series of CDD has a well-defined cyclic pattern with no trend, and another one occurs at Picota (30K.02) where the duration of dry spells has a decreasing trend during the last three decades of the twentieth century. The TAE series of FDD show a different temporal pattern. The frequency of dry days decreases until the mid 1960s, and then has a parabolic behaviour by increasing until the middle of the 1980s and decreasing afterwards. Those findings led us to perform a regional correlation analysis between the CDD and the FDD indices by decade. The Spearman correlation coefficients between these indices were negative but extremely weak as reported before, but this time with one exception: in the last decade of the twentieth century the correlation dropped to  $-0.66$ . These results seem to indicate that, in recent times, an increase in the length of the greatest dry spell entails a decrease in the mean frequency of dry events. Moreover, those dryness indices remain uncorrelated with any of the other indices within that decade.

The TAE series of R5D clearly reflect a strong increase in the short-term precipitation intensity during the last three decades of the twentieth century, except at Lisbon (675) and Aljustrel (26I.03). The regional-average TAE series of SDII, characterizing the precipitation intensity on wet days, shows a cyclic pattern with a small positive tendency after the 1970s. A closer look at the individual stations' TAE series reveals an opposite behaviour at six stations (codes: 675, 25G.01, 25L.01, 25O.01, 26I.03, 27I.01) located in the centre of the study region. These stations show no trend or a small negative tendency during the last three decades of the twentieth century, whereas all other stations have increasing trends in rainfall amounts on wet days. The temporal pattern in the frequency of extremely heavy precipitation events, characterized by the regional-average TAE series of R30, is similar to the pattern of the regional-average TAE series of SDII until the 1970s. Afterwards, the frequency of extreme rainfall has a cyclic pattern with a positive tendency until the mid 1990s and a sudden decrease at the end of this decade. A few stations' TAE series of R30 (stations' codes: 25I.01, 25O.01, 26I.03, 27I.01) exhibit no trend or a small negative tendency in the last three decades of the twentieth century, whereas all other stations have increasing trends in the frequency of heavy rainfall.

The abrupt changes in the trends of CDD in the 1980s, and of SDII and R30 in the 1970s (R5D in less degree), may be partially related to the tendency for the accumulation of positive modes of the North Atlantic Oscillation (NAO) in winter during the last three decades of the twentieth century, which is evident since the 1970s and strengthened during the 1980s and 1990s (Goodess and Jones, 2002; D unkeloh and Jacobeit, 2003; Scaife *et al.*, 2008). A canonical correlation analysis of the CDD index and the frequency of very heavy precipitation days (R90p index) with mean sea-level pressure has revealed that the NAO is an important influence on extreme rainfall over Europe and that the observed trends in these two indices are mainly due to changes in the

NAO (Haylock and Goodess, 2004). The NAO-rainfall relationships tend to be stronger during the wet seasons of the last decades of the twentieth century in southern Portugal (Goodess and Jones, 2002; Trigo *et al.*, 2004), although the NAO impact on winter precipitation is non-stationary, with relatively large oscillations (Trigo *et al.*, 2004). The south-western stations of the Iberia peninsula, including the Serpa and Rel iquias stations in the South of Portugal, have significant correlations between the NAO index and the proportion of rain falling on days with rainfall above the 95th percentile during winter (Rodrigo and Trigo, 2007). However, it is necessary to search for other mechanisms for those stations not linked to NAO (Rodrigo and Trigo, 2007), and further research on the possible causes of the temporal dynamics revealed is required.

The TVE series (moving 5 and 10 years standard deviation statistics) allow the characterization of the extreme precipitation variability through time, and constitute indicators of uncertainty associated with the temporal patterns of the extreme precipitation indices in the period 1955/1999. Figure 7 shows the results of the regional-average TVE series of the six indices. The TVE series of the wetness indices reveal an increase of variability in rainfall frequency and intensity along the 1955/1999 period. The variability of the length of dry spells, characterized by the TVE series of the CDD index, has also increased through time, whereas the variability of the frequency of dry spells (FDD) shows a downward pattern with an increase during the 1990s. These results also support the difficulties with capturing trend signals for these indices at the majority of stations. The TVE series of the AII index reflects a decrease in the dryness variability except during the 1990s. These results indicate that extreme precipitation variability and climate uncertainty are greater in recent times.

#### 4. Conclusion

Whenever the monitoring station network is dense enough, relative approaches are usually preferred to detect inhomogeneities in climate data, because such techniques account for regional climate change and isolate the effects of irregularities in a candidate station by using data from reference stations. The methodology used for the homogeneity assessment of the Portuguese precipitation time series comprised both absolute and relative approaches. During the relative testing stage, we proposed a new technique named *SUR+Ellipse test* that has the advantage of testing simultaneously several candidate series from the same climatic area, taking into account the contemporaneous relationship between them. The results indicate this technique as a valuable tool for homogeneity testing climate time series when the station network is dense enough.

This study confirmed that using absolute approaches without metadata information makes it difficult to determine if changes or lack of changes in a station's time






Figure 7. Ordinary least squares fitting (OLS, thick line) and weighted local polynomial fitting (LOWESS smoothing, thin line) for each regional-average TVE series (moving standard deviation of the extreme index using a time span of 10 years), for the period 1955/99. This figure is available in colour online at [www.interscience.wiley.com/joc](http://www.interscience.wiley.com/joc)

series result from inhomogeneities or simply from abrupt changes in the regional climate. The results of the different procedures implemented were used to develop an overall classification of the daily series using four classes: 'useful', 'potentially useful', 'potentially suspect' and 'suspect'. The intermediate classes were established for stations that were just tested through absolute techniques, since it was not possible to find historic metadata support for the irregularities identified, and testing the 107 series through relative techniques was out of the scope of this work. Therefore, we strongly recommend that further efforts should be made to quality control those series.

From the set of 107 stations compiled for homogeneity assessment, 15 stations with homogeneous daily

records in the period 1955/1999 were selected for temporal patterns analysis. Six precipitation indices were then developed to investigate yearly trends and climate dynamics at the local scale in the South of Portugal. The three dryness indices (AII, CDD and FDD) and the three wetness indices (SDII, R5D and R30) describe moderate climate extremes which are relevant for the management of water resources and land use, modelling of erosion, and other applications for ecosystem and hydrological impact modelling.

The developed AII index is particularly useful to provide information about land vulnerability, especially in agricultural areas such as those located at the South of Portugal. The significant trends of this indicator

in many stations and in the TAE series confirms the desertification-related scenarios that reveal a tendency towards drier climatic conditions in the South of Portugal. The results of the other two dryness indices provide evidence that, in recent times, an increase in the length of the greatest dry spell entails a decrease in the mean frequency of dry events.

The trend signals of the wetness indices were not significant at the majority of stations. However, the moving window techniques revealed an increase in the short-term precipitation intensity (R5D index) during the last three decades of the twentieth century. Indices characterizing the precipitation intensity on wet days (SDII) and the frequency of extremely heavy precipitation events (R30) have cyclic patterns and different trend signals at the local scale, during the period 1955/1999. Finally, the results also indicate that extreme precipitation variability and climate uncertainty are greater in recent times.

### Acknowledgements

This study was partially supported by the Fundação para a Ciência e Tecnologia (FCT) as a part of the project CIDmeg – Construction of a Desertification Susceptibility Index for the Left Margin of Guadiana, contract no. POCI/CLI/58865/2004. The authors also thank two anonymous reviewers for valuable suggestions, which helped them to improve the quality of this paper.

### A1. Appendix

#### A1.1. SUR+Ellipse test

Zellner (1962) proposed the SUR approach for situations where at least two equations are being estimated and the error terms are contemporaneously but not serially correlated. In a general specification of  $M$  seemingly unrelated regression equations (one for each candidate series), the  $i$ th equation is given by

$$Y_i = X_i \beta_i + e_i, \quad i = 1, \dots, M \quad (\text{A1})$$

where  $Y_i$  is a vector of dimension  $(T \times 1)$  containing all the observations on the  $i$ th dependent variable ( $T$  observations from the  $i$ th candidate series);  $X_i$  is a matrix of dimension  $(T \times K_i)$  containing all the observations from all the  $K_i$  explanatory variables ( $T$  observations from each reference series, and  $T$  observations from the constant term);  $\beta_i$  is a vector, of dimension  $(K_i \times 1)$ , of unknown coefficients to be estimated;  $e_i$  is a vector of dimension  $(T \times 1)$  containing the error terms for all observations. Note that each equation involves  $K_i$  regressors, meaning that each equation need not have the same number of explanatory variables. However, if all equations have identical explanatory variables, then generalized least squares are equivalent to equation-by-equation OLS (Greene, 2003; p. 343). Thus, for those situations the proposed approach is equivalent to the

method of cumulative residuals (*Ellipse test*) described by Allen *et al.* (1998). Combining all equations into one model yields

$$\begin{bmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_M \end{bmatrix} = \begin{bmatrix} X_1 & 0 & \cdots & 0 \\ 0 & X_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & X_M \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_M \end{bmatrix} + \begin{bmatrix} e_1 \\ e_2 \\ \vdots \\ e_M \end{bmatrix} \quad (\text{A2})$$

Or, alternatively,  $Y = X\beta + e$ , where the definitions of  $Y$ ,  $X$ ,  $\beta$  and  $e$  are obvious from Equation (A2) and their dimensions are, respectively,  $(MT \times 1)$ ,  $(MT \times K)$ ,  $(K \times 1)$  and  $(MT \times 1)$ , with  $K = \sum_{i=1}^M K_i$ . Given that  $e_{it}$  is the error for the  $i$ th equation in the  $t$ th time period, the assumption of contemporaneous disturbance correlation, but not correlation over time, implies that the covariance matrix for the complete error vector can be written as

$$W = E[ee'] = \Sigma \otimes I_T, \quad \Sigma = \begin{bmatrix} \sigma_{11} & \sigma_{12} & \cdots & \sigma_{1M} \\ \sigma_{21} & \sigma_{22} & \cdots & \sigma_{2M} \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{M1} & \sigma_{M2} & \cdots & \sigma_{MM} \end{bmatrix} \quad (\text{A3})$$

where  $\otimes$  denotes the Kronecker product, indicating that each element of  $\Sigma$  is multiplied by an identity matrix. The matrix  $\Sigma$  is symmetric, so that  $\sigma_{ij} = \sigma_{ji}$  and it is non-singular, and thus has an inverse. The *generalized least squares estimator* of  $\beta$ , typically denoted by  $\hat{\beta}$ , is the best linear unbiased estimator, but assumes that  $\Sigma$  is known. In practice, the variances and covariances are usually unknown and must be estimated, thus the *feasible generalized least squares estimator*  $\hat{\hat{\beta}}$  is generally used (Greene, 2003; p. 344), and  $\Sigma$  is replaced by a consistent estimator. To estimate the  $\sigma_{ij}$ , each equation is first estimated by OLS in order to obtain the least squares residuals  $\hat{e}_i$ . Consistent estimates of the variances and covariances are then given by

$$\hat{\sigma}_{ij} = \frac{1}{T} \hat{e}_i' \hat{e}_j = \frac{1}{T} \sum_{t=1}^T \hat{e}_{it} \hat{e}_{jt} \quad (\text{A4})$$

If we define  $\hat{\Sigma}$  as the matrix  $\Sigma$  with the unknown  $\sigma_{ij}$  replaced by  $\hat{\sigma}_{ij}$ , then the feasible generalized least squares estimator for  $\beta$  can be written as

$$\hat{\hat{\beta}} = \left[ X' \left( \hat{\Sigma}^{-1} \otimes I_T \right) X \right]^{-1} X' \left( \hat{\Sigma}^{-1} \otimes I_T \right) Y \quad (\text{A5})$$

The SUR + Ellipse test uses the cumulative residuals from such a SUR model to identify inhomogeneities in several candidate series from the same climatic area. A candidate series can be considered homogeneous if the cumulative residuals are not biased. The bias hypothesis

can be tested using an ellipse defining the confidence limits (Allen *et al.*, 1998). For each equation  $i$  ( $i$ th candidate series), the axes and the parametric equation of the  $i$ th ellipse are, respectively:

$$\begin{cases} \alpha_i = T/2 \\ \beta_i = \frac{T}{\sqrt{T-1}} z_p S_{e,i} \end{cases}, \quad \begin{cases} X_i = \alpha_i \cos(\theta) \\ Y_i = \beta_i \sin(\theta) \end{cases} \quad (\text{A6})$$

where  $z_p$  stands for the standard normal variate for the desired probability  $p$  (confidence level);  $S_{e,i}$  is the standard deviation of the residuals of the  $i$ th equation; and  $\theta$  [rad] varies from 0 to  $2\pi$ . Plotting the cumulative residuals against time, using the time scale (interval) of the variable under analysis, the accumulated residual curve is obtained. If all the cumulative residuals lie inside the ellipse then the hypothesis of homogeneity is not rejected for the significance level considered.

## References

- Aguilar E, Auer I, Brunet M, Peterson TC, Wieringa J. 2003. *Guidelines on Climate Metadata and Homogenization*. World Meteorological Organization: WMO-TD No. 1186, WCDMP No. 53: Switzerland, Geneva.
- Alexandersson H. 1986. A homogeneity test applied to precipitation data. *Journal of Climatology* **6**: 661–675, DOI: 10.1002/joc.3370060607.
- Alexandersson H, Moberg A. 1997. Homogenization of Swedish temperature data. Part I: Homogeneity test for linear trends. *International Journal of Climatology* **17**: 25–34, DOI: 10.1002/(SICI)1097-0088(199701)17:1<25::AID-JOC103>3.0.CO;2-J.
- Allen RG, Pereira LS, Raes D, Smith M. 1998. Statistical analysis of weather data sets. In *Crop Evapotranspiration – Guidelines for Computing Crop Water Requirements*, FAO – Food and Agriculture Organization of the United Nations (ed.). FAO Irrigation and drainage paper 56, Annex IV, Rome.
- Auer I, Böhm R, Jurković A, Orlik A, Potzmann R, Schöner W, Ungersböck M, Brunetti M, Nanni T, Maugeri M, Briffa K, Jones P, Efthymiadis D, Mestre O, Moisselin J-M, Begert M, Brazdil R, Bochnicek O, Cegnar T, Gajić-Eapka M, Zaninović K, Majstorović Ž, Szalai S, Szentimrey T, Mercalli L. 2005. A new instrumental precipitation dataset for the greater alpine region for the period 1800–2002. *International Journal of Climatology* **25**: 139–166, DOI: 10.1002/joc.1135.
- Buishand TA. 1982. Some methods for testing the homogeneity of rainfall records. *Journal of Hydrology* **58**: 11–27, DOI: 10.1016/0022-1694(82)90066-X.
- Ceballos A, Martínez-Fernández J, Luengo-Ugidos JA. 2004. Analysis of rainfall trends and dry periods on a pluviometric gradient representative of Mediterranean climate in the Duero Basin, Spain. *Journal of Arid Environments* **58**: 215–233, DOI: 10.1016/j.jaridenv.2003.07.002.
- Cleveland WS. 1979. Robust locally weighted regression and smoothing scatterplots. *Journal of the American Statistical Association* **74**: 829–836, DOI: 10.2307/2286407.
- Costa AC, Negreiros J, Soares A. 2008. Identification of inhomogeneities in precipitation time series using stochastic simulation. In *geoENV VI – Geostatistics for Environmental Applications*, Soares A, Pereira MJ, Dimitrakopoulos R (eds). Springer: Netherlands; 275–282.
- Ducré-Robitaille JF, Vincent LA, Boulet G. 2003. Comparison of techniques for detection of discontinuities in temperature series. *International Journal of Climatology* **23**: 1087–1101, DOI: 10.1002/joc.924.
- Dünkeloh A, Jacobeit J. 2003. Circulation dynamics of Mediterranean precipitation variability 1948–1998. *International Journal of Climatology* **23**: 1843–1866, DOI: 10.1002/joc.973.
- Engle RF. 1982. Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. *Econometrica* **50**: 987–1007, DOI: 10.2307/1912773.
- Feng S, Hu Q, Qian W. 2004. Quality control of daily meteorological data in China, 1951–2000: a new dataset. *International Journal of Climatology* **24**: 853–870, DOI: 10.1002/joc.1047.
- Frich P, Alexander LV, Della-Marta P, Gleason B, Haylock M, Klein Tank AMG, Peterson T. 2002. Observed coherent changes in climatic extremes during the second half of the twentieth century. *Climate Research* **19**: 193–212, DOI: 10.3354/cr019193.
- Goodess CM, Jones PD. 2002. Links between circulation and changes in the characteristics of Iberian rainfall. *International Journal of Climatology* **22**: 1593–1615, DOI: 10.1002/joc.810.
- Greene WH. 2003. *Econometric Analysis*, 5th edn. Prentice Hall International Edition, Pearson Education: New Jersey.
- Haylock MR, Goodess CM. 2004. Interannual variability of European extreme winter rainfall and links with mean large-scale circulation. *International Journal of Climatology* **24**: 759–776, DOI: 10.1002/joc.1033.
- Haylock M, Nicholls N. 2000. Trends in extreme rainfall indices for an updated high quality data set for Australia, 1910–1998. *International Journal of Climatology* **20**: 1533–1541, DOI: 10.1002/1097-0088(20001115)20:13<1533::AID-JOC586>3.0.CO;2-J.
- Kendall MG. 1975. *Rank Correlation Methods*. Charles Griffin: London.
- Kiktev D, Sexton DMH, Alexander L, Folland CK. 2003. Comparison of modeled and observed trends in indices of daily climate extremes. *Journal of Climate* **16**: 3560–3571, DOI: 10.1175/1520-0442(2003)016<3560:COMAOT>2.0.CO;2.
- Klein Tank AMG, Können GP. 2003. Trends in indices of daily temperature and precipitation extremes in Europe, 1946–1999. *Journal of Climate* **16**: 3665–3680, DOI: 10.1175/1520-0442(2003)016<3665:TIHODT>2.0.CO;2.
- Klein Tank AMG, Wijngaard JB, Können GP, Böhm R, Demarée G, Gocheva A, Mileta M, Pashiardis S, Hejkrlik L, Kern-Hansen C, Heino R, Bessemoulin P, Müller-Westermeier G, Tzanakou M, Szalai S, Pálsdóttir T, Fitzgerald D, Rubin S, Capaldo M, Maugeri M, Leitass A, Bukantis A, Aberfeld R, van Engelen AFV, Forland E, Miettus M, Coelho F, Mares C, Razuvaev V, Nieplova E, Cegnar T, López JA, Dahlström B, Moberg A, Kirchhofer W, Ceylan A, Pachaliuk O, Alexander LV, Petrovic P. 2002. Daily dataset of 20th-century surface air temperature and precipitation series for the European Climate Assessment. *International Journal of Climatology* **22**: 1441–1453, DOI: 10.1002/joc.773.
- Kostopoulou E, Jones PD. 2005. Assessment of climate extremes in the Eastern Mediterranean. *Meteorology and Atmospheric Physics* **89**: 69–85, DOI: 10.1007/s00703-005-0122-2.
- Lana X, Martínez MD, Burgueño A, Serra C, Martín-Vide J, Gómez L. 2008. Spatial and temporal patterns of dry spell lengths in the Iberian Peninsula for the second half of the twentieth century. *Theoretical and Applied Climatology* **91**: 99–116, DOI: 10.1007/s00704-007-0300-x.
- Lanzante JR. 1996. Resistant, robust and nonparametric techniques for the analysis of climate data: Theory and examples, including applications to historical radiosonde station data. *International Journal of Climatology* **16**: 1197–1226, DOI: 10.1002/(SICI)1097-0088(199611)16:11<1197::AID-JOC89>3.0.CO;2-L.
- Mann HB. 1945. Non-parametric test against trend. *Econometrica* **13**: 245–259, DOI: 10.2307/1907187.
- Moberg A, Jones PD. 2005. Trends in indices for extremes in daily temperature and precipitation in central and western Europe, 1901–1999. *International Journal of Climatology* **25**: 1149–1171, DOI: 10.1002/joc.1163.
- Nicolau RC. 1999. *Avaliação da qualidade de séries de registos de precipitação anual com vista à seleção, ajustamento e completamento das mesmas para posterior modelação da variabilidade espacial da pluviosidade*. Centro Nacional de Investigação Geográfica, Estudo, 053C/19, January 1999, CNIG, Lisboa.
- Peterson TC, Easterling DR, Karl TR, Groisman P, Nicholls N, Plummer N, Torok S, Auer I, Boehm R, Gullett D, Vincent L, Heino R, Tuomenvirta H, Mestre O, Szentimrey T, Salinger J, Forland EJ, Hansen-Bauer I, Alexandersson H, Jones P, Parker D. 1998. Homogeneity adjustments of in situ atmospheric climate data: A review. *International Journal of Climatology* **18**: 1493–1517, DOI: 10.1002/(SICI)1097-0088(19981115)18:13<1493::AID-JOC329>3.0.CO;2-T.
- Peterson TC, Folland C, Gruza G, Hogg W, Mokssit A, Plummer N. 2001. *Report on the Activities of the Working Group on Climate Change Detection and Related Rapporteurs 1998–2001*. World Meteorological Organization, WCDMP–No. 47/WMO–TD No. 1071: Geneva, Switzerland.

- Pettit AN. 1979. A non-parametric approach to the change-point detection. *Applied Statistics* **28**: 126–135.
- Reeves J, Chen J, Wang XL, Lund R, Lu Q. 2007. A review and comparison of change-point detection techniques for climate data. *Journal of Applied Meteorology and Climatology* **46**: 900–915, DOI: 10.1175/JAM2493.1.
- Rodrigo FS, Trigo RM. 2007. Trends in daily rainfall in the Iberian Peninsula from 1951 to 2002. *International Journal of Climatology* **27**: 513–529, DOI: 10.1002/joc.1409.
- Rousseeuw PJ, Croux C. 1993. Alternatives to the median absolute deviation. *Journal of the American Statistical Association* **88**: 1273–1283, DOI: 10.2307/2291267.
- SAS Institute. 1999. *SAS Procedures Guide*, Version 8. SAS Institute Inc.: Cary, NC.
- Scaife AA, Folland CK, Alexander LV, Moberg A, Knight JR. 2008. European climate extremes and the North Atlantic Oscillation. *Journal of Climate* **21**: 72–83, DOI: 10.1175/2007JCLI1631.1.
- Soares A. 2001. Direct sequential simulation and cosimulation. *Mathematical Geology* **33**: 911–926, DOI: 10.1023/A:1012246006212.
- Trigo RM, DaCamara C. 2000. Circulation weather types and their impact on the precipitation regime in Portugal. *International Journal of Climatology* **20**: 1559–1581, DOI: 10.1002/1097-0088(20001115)20:13<1559::AID-JOC555>3.0.CO;2-5.
- Trigo RM, Pozo-Vázquez D, Osborn TJ, Castro-Díez Y, Gámiz-Fortis S, Esteban-Parra MJ. 2004. North Atlantic Oscillation influence on precipitation, river flow and water resources in the Iberian Peninsula. *International Journal of Climatology* **24**: 925–944, DOI: 10.1002/joc.1048.
- Tuomenvirta H. 2001. Homogeneity adjustments of temperature and precipitation series – Finnish and Nordic data. *International Journal of Climatology* **21**: 495–506, DOI: 10.1002/joc.616.
- Vicente-Serrano SM, Cuadrat-Prats JM. 2007. Trends in drought intensity and variability in the middle Ebro valley (NE of the Iberian peninsula) during the second half of the twentieth century. *Theoretical and Applied Climatology* **88**: 247–258, DOI: 10.1007/s00704-006-0236-6.
- Von Neumann J. 1941. Distribution of the ratio of the mean square successive difference to the variance. *Annals of Mathematical Statistics* **13**: 367–395.
- Wald A, Wolfowitz J. 1943. An exact test for randomness in the non-parametric case based on serial correlation. *Annals of Mathematical Statistics* **14**: 378–388.
- Wijngaard JB, Klein Tank AMG, Können GP. 2003. Homogeneity of 20th century European daily temperature and precipitation series. *International Journal of Climatology* **23**: 679–692, DOI: 10.1002/joc.906.
- Zellner A. 1962. An efficient method of estimating seemingly unrelated regressions and tests for aggregation bias. *Journal of the American Statistical Association* **57**: 348–368, DOI: 10.2307/2281644.